

CUSHING'S DISEASE IN HORSES

James M. Casey, D.V.M, M.S.
Equine Sports Medicine, Dentistry, & Surgery

Office: (301) 725-4371 Cell: (301) 785-5167
Website: www.equinehorsevet.com

PITUITARY PARS INTERMEDIA DYSFUNCTION (EQUINE CUSHING'S DISEASE)

Pituitary *pars intermedia* dysfunction (PPID, previously known as Equine Cushing's Disease) is a complex condition associated with abnormal function of a small, hormone-producing organ, the pituitary gland, which lies at the base of the brain. The cause is not fully known but currently it is thought that as part of the ageing process, some horses develop enlargement of part of the pituitary gland (the pars intermedia), which then produces excessive amounts of a range of hormones important in controlling the body. The hormones in question are known as the pro-opiomelanocortin-derived peptides (POMC) and include melanocytstimulating hormone (MSH), corticotropin-like intermediate lobe peptide (CLIP), beta-endorphin (β -END) and adrenocorticotrophic hormone (ACTH). The fact that there are several hormones involved is the reason that the name Equine Cushing's disease has been abandoned in favour of PPID since in other species Cushing's disease generally involves only one hormone system. The POMC hormones play a part in controlling a wide range of body functions and the hormonal imbalances caused in PPID result in a range of clinical abnormalities.



Excessively long hair coat in old horse with PPID

What are the clinical signs?

The average age of a horse or pony with PPID is 20 years, although horses as young as 14 years can be affected. The most striking sign, but one which is not always present, especially in early cases, is an excessively long and curly coat (hirsutism). This is caused by the hair coat not being shed normally. This 'woolly bear' appearance may have been preceded by several years during which the winter hair coat was not shed until very late. Many affected horses drink excessive amounts of water and produce large volumes of urine (polydipsia and polyuria). Other signs are lethargy, weight loss or an inability to maintain muscle while accumulating fat (especially on the crest and on top of the rump) and excessive sweating. Affected cases often suffer from acute or recurrent attacks of laminitis and many suffer from liver disease. PPID can be severely debilitating and affected horses or ponies can develop severe pneumonia, dental infections or other secondary infections.


How is the condition diagnosed?

In aged horses or ponies with hair coats that are several inches long and curly and there is a history of laminitis and/or excessive drinking, the diagnosis can often be made on these grounds alone. In other less obvious cases, especially in younger horses that are developing the condition, the diagnosis depends on a combination of clinical signs, blood and urine tests. Due to the complex nature of the condition, it may be necessary to use additional specialized blood tests which look at changes in hormone levels in response to the injection of other hormones, testing hormone system function. Animals with PPID usually respond abnormally to one or more of these tests, but interpretation of results is not always straight forward, as they can sometimes appear contradictory. Your veterinarian will help you with this.

Is any treatment available?

Due to the complex, multi-organ problems that can be associated with PPID, attempts at treatment must include attention to all aspects of the animal's condition. Affected horses are more susceptible to parasites and infection so regular worming, vaccination and teeth care should be instituted. Any dietary changes will depend on the age and condition of the horse or pony, the available grazing, the presence or absence of laminitis, expectations for work, etc and should be discussed with your veterinarian. Cases with excessively long coats may sweat excessively or may take a long time to dry off after exercise and they should be clipped over their body and neck to help them dry off more quickly. In cooler weather, clipped horses and ponies may need to have a blanket in very cold weather.

There are three medicines that are now used to treat PPID in horses and ponies. These are called pergolide mesylate, trilostane and cyproheptadine. None of these drugs is totally effective in all cases and some cases do not respond to treatment at all. Treatment with either must continue for the life of the horse or pony and is not inexpensive. Treated cases must be regularly monitored with regard to their clinical signs and sometimes by repeat blood tests. Your veterinarian is the best person to discuss this complex problem, to determine whether or not your horse or pony really does have PPID and if so what treatment is appropriate.



Enlarged pituitary gland containing tumor (adenoma) from horse with PPID

Can it be prevented?

It is not possible to prevent this condition. However, with available blood tests we are now able to recognize and treat more cases earlier and more effectively and many treated horses and ponies go on to live normal lives for many years after diagnosis and treatment.

Equine Sports Medicine, Dentistry, & Surgery
James M. Casey, D.V.M, M.S.
Veterinary Practitioner and Surgeon

Practicing in the Mid-Atlantic States & Florida
(MD, VA, DE, PA, NJ, GA, KY, FL)

Office: 301-725-4371

Cell: 301-785-5167

Email: casey@us.net

Website: www.equinehorsevet.com

Consults Nationally and Worldwide